arXiv:0905.2024v1 [math.AP] 13 May 2009

BOUNDARY-VALUE PROBLEMS WITH NON-LOCAL INITIAL CONDITION FOR PARABOLIC EQUATIONS WITH PARAMETER

J.M.Rassias[1] **and E.T.Karimov**[2]

[1]*Pedagogical department, Mathematics and Informatics section, National and Capodostrian University of Athens. jrassias@primedu.uoa.gr*

[2]*Institute of Mathematics and Information Technologies, Uzbek Academy of Sciences. E-mail: erkinjon@gmail.com*

Abstract

In 2002, J.M.Rassias (Uniqueness of quasi-regular solutions for bi-parabolic elliptic bi-hyperbolic Tricomi problem, Complex Variables, 47 (8) (2002), 707-718) imposed and investigated the bi-parabolic elliptic bi-hyperbolic mixed type partial differential equation of second order. In the present paper some boundary-value problems with non-local initial condition for model and degenerate parabolic equations with parameter were considered. Also uniqueness theorems are proved and non-trivial solutions of certain non-local problems for forward-backward parabolic equation with parameter are investigated at specific values of this parameter by employing the classical "a-b-c" method. Classical references in this field of mixed type partial differential equations are given by: J.M.Rassias (Lecture Notes on Mixed Type Partial Differential Equations, World Scientific, 1990, pp.1-144) and M.M.Smirnov (Equations of Mixed Type, Translations of Mathematical Monographies, 51, American Mathematical Society, Providence, R.I., 1978 pp.1-232). Other investigations are achieved by G.C.Wen et al. (in period 1990-2007).

MSC 2000 classification:

Keywords: Degenerate parabolic equation; forward-backward parabolic equation with a parameter; boundary-value problems with non-local initial conditions; classical "a-b-c" method.

1 Introduction

Degenerate partial differential equations have numerous applications in Aerodynamics and Hydrodynamics. For example, problems for mixed subsonic and supersonic flows were considered by F.I.Frankl [1]. Reviews of interesting results on degenerated elliptic and hyperbolic equations up to 1965, one can find in the book by M.M.Smirnov [2]. Among other research results on this kind of equations were investigated by J.M.Rassias [3-10], G.C.Wen [11-15], A.Hasanov [16] and references therein. Also works by M.Gevrey [17], A.Friedman [18], Yu.Gorkov [19] are well-known on construction fundamental solutions for degenerated parabolic equations. In addition it was studied by N.N.Shopolov [20] a boundary-value problem with initial non-local condition for model parabolic equation in [21-25]. However, in 2002, J.M.Rassias (Uniqueness of quasi-regular solutions for bi-parabolic elliptic bi-hyperbolic Tricomi problem, Complex Variables, 47 (8) (2002), 707-718) imposed and investigated the bi-parabolic elliptic bi-hyperbolic mixed type partial differential equation of second order. In the present paper some boundary-value problems with non-local initial condition for model and degenerate parabolic equations with parameter were considered. Also uniqueness theorems are proved and non-trivial solutions of certain non-local problems for forward-backward parabolic equation with parameter are investigated at specific values of this parameter by employing the classical "a-b-c" method. Classical references in this field of mixed type partial differential equations are given by: J.M.Rassias (Lecture Notes on Mixed Type Partial Differential Equations, World Scientific, 1990, pp.1-144) and M.M.Smirnov (Equations of Mixed Type, Translations of Mathematical Monographies, 51, American Mathematical Society, Providence, R.I., 1978 pp.1-232). Other investigations are achieved by G.C.Wen et al. (in period 1990-2007).

2 Non-local problems for degenerate parabolic equations with parameter.

Let us consider a parabolic equation

$$y^m u_{xx} - x^n u_y - \lambda x^n y^m u = 0, \tag{1}$$

with two lines of degeneration in the domain $\Phi = \{(x, y) : \ 0 < x < 1, \ 0 < y < 1\}$, where $m, \ n > 0, \ \lambda \in C$.

The problem 1. To find a regular solution of the equation satisfying boundary conditions

$$u\left(0, y\right) = 0, \quad u\left(1, y\right) = 0, \quad 0 \leq y \leq 1, \tag{2}$$

and non-local initial condition

$$u\left(x, 0\right) = \alpha u\left(x, 1\right), \ 0 \leq x \leq 1, \tag{3}$$

where α is non-zero real number.

The following statements are true:

Theorem 1. Let $\alpha \in [-1, 0) \cup (0, 1]$, $\mathrm{Re}\lambda \geq 0$. If there exists a solution of the problem 1, then it is unique.

Corollary 1. The problem 1 can have non-trivial solutions only when parameter λ lies outside of the sector $\Delta = \{\lambda : \ \mathrm{Re}\lambda \geq 0\}$. These non-trivial solutions represented by

$$u_{pk}\left(x, y\right) = C_{pk} \left(\frac{2}{n+2}\right)^{\frac{1}{n+2}} \mu_k^{\frac{1}{2(n+2)}} x^{\frac{1}{2}} I_{\frac{1}{n+2}} \left(\frac{2\sqrt{\mu_k}}{n+2} x^{\frac{n+2}{2}}\right) e^{(-\ln|\alpha| - ip\pi) y^{m+1}}, \tag{4}$$

where C_{pk} are constants, p, k are real positive numbers. Eigenvalues defined as

$$\lambda_{pk} = \mu_k + (m+1) \ln|\alpha| + i\left(m+1\right) p\pi.$$

Here μ_k are roots of the equation

$$I_{\frac{1}{n+2}} \left(\frac{2\sqrt{\mu}}{n+2}\right) = 0,$$

where $I_s\left(\right)$ is the first kind modified Bessel function of s-th order.

We will omit the proof, because further we consider similar problem in three-dimensional domain in a full detail.

Let Ω be a simple-connected bounded domain in R^3 with boundaries S_i $(i=\overline{1,6})$. Here

$$S_1=\{(x,y,t):\ t=0,\ 0<x<1,\ 0<y<1\}\,,\ S_2=\{(x,y,t):\ x=1,\ 0<y<1,\ 0<t<1\}\,,$$
$$S_3=\{(x,y,t):\ y=0,\ 0<x<1,\ 0<t<1\}\,,\ S_4=\{(x,y,t):\ x=0,\ 0<y<1,\ 0<t<1\}\,,$$
$$S_5=\{(x,y,t):\ y=1,\ 0<x<1,\ 0<t<1\}\,,\ S_6=\{(x,y,t):\ t=1,\ 0<x<1,\ 0<y<1\}\,.$$

We consider the following degenerate parabolic equation

$$x^n y^m u_t = y^m u_{xx} + x^n u_{yy} - \lambda x^n y^m u \tag{5}$$

in the domain Ω. Here $m>0$, $n>0$, $\lambda=\lambda_1+i\lambda_2$, $\lambda_1,\lambda_2\in R$.

The problem 2. To find a function $u\,(x,y,t)$ satisfying the following conditions:

i) $u\,(x,y,t)\in C\left(\overline{\Omega}\right)\cap C^{2,2,1}_{x,y,t}\,(\Omega)$;

ii) $u\,(x,y,t)$ satisfies the equation (5) in Ω;

iii) $u\,(x,y,t)$ satisfies boundary conditions

$$u\,(x,y,t)\,|_{S_2\cup S_3\cup S_4\cup S_5}=0\ ; \tag{6}$$

iv) and non-local initial condition

$$u\,(x,y,0)=\alpha u\,(x,y,1)\,. \tag{7}$$

Here $\alpha=\alpha_1+i\alpha_2$, α_1,α_2 are real numbers, moreover $\alpha_1^2+\alpha_2^2\neq 0$.

Theorem 2. If $\alpha_1^2+\alpha_2^2<1$, $\lambda_1\geq 0$ and exists a solution of the problem 2, then it is unique.

Proof:

Let us suppose that the problem 2 has two u_1, u_2 solutions. Denoting $u=u_1-u_2$ we claim that $u\equiv 0$ in Ω.

First we multiply equation (5) to the function $\overline{u}\,(x,y,t)$, which is complex conjugate function of $u\,(x,y,t)$. Then integrate it along the domain Ω_ε with boundaries

$$S_{1\varepsilon}=\{(x,y,t):\ t=\varepsilon,\ \varepsilon<x<1-\varepsilon,\ \varepsilon<y<1-\varepsilon\}\,,$$
$$S_{2\varepsilon}=\{(x,y,t):\ x=1-\varepsilon,\ \varepsilon<y<1-\varepsilon,\ \varepsilon<t<1-\varepsilon\}\,,$$
$$S_{3\varepsilon}=\{(x,y,t):\ y=\varepsilon,\ \varepsilon<x<1-\varepsilon,\ \varepsilon<t<1-\varepsilon\}\,,$$
$$S_{4\varepsilon}=\{(x,y,t):\ x=\varepsilon,\ \varepsilon<y<1-\varepsilon,\ \varepsilon<t<1-\varepsilon\}\,,$$
$$S_{5\varepsilon}=\{(x,y,t):\ y=1-\varepsilon,\ \varepsilon<x<1-\varepsilon,\ \varepsilon<t<1-\varepsilon\}\,,$$
$$S_{6\varepsilon}=\{(x,y,t):\ t=1-\varepsilon,\ \varepsilon<x<1-\varepsilon,\ \varepsilon<y<1-\varepsilon\}\,.$$

Then taking real part of the obtained equality and considering

$$\mathrm{Re}\,(y^m\overline{u}u_{xx})=\mathrm{Re}\,(y^m\overline{u}u_x)_x-y^m\left|u_x\right|^2,\ \mathrm{Re}\,(x^n\overline{u}u_{yy})=\mathrm{Re}\,(x^n\overline{u}u_y)_y-x^n\left|u_y\right|^2,$$

$$\mathrm{Re}\,(x^n y^m\overline{u}u_t)=\left(\frac{1}{2}x^n y^m\left|u\right|^2\right)_t,$$

after using Green's formula we pass to the limit at $\varepsilon \to 0$. Then we get

$$\int\limits_{\partial\Omega}\int \mathrm{Re}\left[y^m\overline{u}u_x\cos(\nu,x)+x^n\overline{u}u_y\cos(\nu,y)-\tfrac{1}{2}x^ny^m\left|u\right|^2\cos(\nu,t)\right]d\tau$$
$$=\int\int\limits_{\Omega}\int\left(y^m\left|u_x\right|^2+x^n\left|u_y\right|^2+\lambda_1x^ny^m\left|u\right|\right)d\sigma$$

where ν is outer normal. Taking into account $\mathrm{Re}\left[\overline{u}u_x\right]=\mathrm{Re}\left[u\overline{u}_x\right],\ \mathrm{Re}\left[\overline{u}u_y\right]=\mathrm{Re}\left[u\overline{u}_y\right]$ we obtain

$$\mathrm{Re}\int\int\limits_{S_1}\frac{1}{2}x^ny^m\left|u\right|^2d\tau_1+\int\int\limits_{S_2}y^m\mathrm{Re}\left[u\overline{u}_x\right]d\tau_2-\int\int\limits_{S_3}x^n\mathrm{Re}\left[u\overline{u}_y\right]d\tau_3-\int\int\limits_{S_4}y^m\mathrm{Re}\left[u\overline{u}_x\right]d\tau_4+$$

$$+\int\int\limits_{S_5}x^n\mathrm{Re}\left[u\overline{u}_y\right]d\tau_5-\mathrm{Re}\int\int\limits_{S_6}\frac{1}{2}x^ny^m\left|u\right|^2d\tau_6=\int\int\limits_{\Omega}\int\left(y^m\left|u_x\right|^2+x^n\left|u_y\right|^2+\lambda_1x^ny^m\left|u\right|\right)d\sigma. \quad (8)$$

From (8) and by using conditions (6), (7), we find

$$\frac{1}{2}\left[1-\left(\alpha_1^2+\alpha_2^2\right)\right]\int\limits_0^1\int\limits_0^1x^ny^m\left|u\left(x,y,1\right)\right|dxdy+\int\int\limits_{\Omega}\int\left(y^m\left|u_x\right|^2+x^n\left|u_y\right|^2+\lambda_1x^ny^m\left|u\right|\right)d\sigma=0. \quad (9)$$

Setting $\alpha_1^2+\alpha_2^2<1,\ \lambda_1\geq 0$, from (9) we have $u\left(x,y,t\right)\equiv 0$ in $\overline{\Omega}$.

Theorem is proved.

We find below non-trivial solutions of the problem 2 at some values of parameter λ for which the uniqueness condition $\mathrm{Re}\lambda=\lambda_1\geq 0$ is not fulfilled.

We search the solution of Problem 2 as follows

$$u\left(x,y,t\right)=X\left(x\right)\cdot Y\left(y\right)\cdot T\left(t\right). \quad (10)$$

After some evaluations we obtain the following eigenvalue problems:

$$\begin{cases} X''\left(x\right)+\mu_1x^nX\left(x\right)=0\\ X\left(0\right)=0,\ \ X\left(1\right)=0;\end{cases} \quad (11)$$

$$\begin{cases} Y''\left(y\right)+\mu_2y^mY\left(y\right)=0\\ Y\left(0\right)=0,\ \ Y\left(1\right)=0;\end{cases} \quad (12)$$

$$\begin{cases} T'\left(t\right)+\left(\lambda+\mu\right)T\left(t\right)=0\\ T\left(0\right)=\alpha T\left(1\right).\end{cases} \quad (13)$$

Here $\mu=\mu_1+\mu_2$ is a Fourier constant.

Solving eigenvalue problems (11), (12) we find

$$\mu_{1k}=\left(\frac{n+2}{2}\widetilde{\mu_{1k}}\right)^2,\ \ \mu_{2p}=\left(\frac{m+2}{2}\widetilde{\mu_{2p}}\right)^2, \quad (14)$$

$$X_k\left(x\right)=A_k\left(\frac{2}{n+2}\right)^{\frac{1}{n+2}}\mu_{1k}^{\frac{1}{2(n+2)}}x^{\frac{1}{2}}J_{\frac{1}{n+2}}\left(\frac{2\sqrt{\mu_{1k}}}{n+2}x^{\frac{n+2}{2}}\right), \quad (15)$$

$$Y_p\left(y\right)=B_p\left(\frac{2}{m+2}\right)^{\frac{1}{m+2}}\mu_{2p}^{\frac{1}{2(m+2)}}y^{\frac{1}{2}}J_{\frac{1}{m+2}}\left(\frac{2\sqrt{\mu_{2p}}}{m+2}x^{\frac{m+2}{2}}\right), \quad (16)$$

where $k,p=1,2,...,$ $\widetilde{\mu_{1k}}$ and $\widetilde{\mu_{2p}}$ are roots of equations $J_{\frac{1}{n+2}}\left(x\right)=0$ and $J_{\frac{1}{m+2}}\left(y\right)=0$, respectively.

The eigenvalue problem (13) has non-trivial solution only when $\begin{cases} \alpha_1 = e^{\lambda_1+\mu_{kp}}\cos\lambda_2 \\ \alpha_2 = e^{\lambda_1+\mu_{kp}}\sin\lambda_2. \end{cases}$

Here $\lambda = \lambda_1 + i\lambda_2,\ \ \alpha = \alpha_1 + i\alpha_2,\ \ \mu_{kp} = \mu_{1k} + \mu_{2p}$. After elementary calculations, we get

$$\lambda_1 = -\mu_{kp} + \ln\sqrt{\alpha_1^2+\alpha_2^2},\ \ \lambda_2 = \arctan\frac{\alpha_2}{\alpha_1} + s\pi,\ \ s \in Z^+ \tag{17}$$

Corresponding eigenfunctions have the form

$$T_{kp}\left(t\right) = C_{kp}e^{\left[\mu_{kp} - \ln\sqrt{\alpha_1^2+\alpha_2^2} - i\left(\arctan\frac{\alpha_2}{\alpha_1}+s\pi\right)\right]t}. \tag{18}$$

Considering (10), (15), (16) and (18) we can write non-trivial solutions of the problem 2 in the following form:

$$u_{kp}\left(x,y,t\right) = D_{kp}\left(\frac{2}{n+2}\right)^{\frac{1}{n+2}}\left(\frac{2}{m+2}\right)^{\frac{1}{m+2}}\mu_{1k}^{\frac{1}{2(n+2)}}\mu_{2p}^{\frac{1}{2(m+2)}}\sqrt{xy}J_{\frac{1}{n+2}}\left(\frac{2\sqrt{\mu_{1k}}}{n+2}x^{\frac{n+2}{2}}\right)$$

$$\times J_{\frac{1}{m+2}}\left(\frac{2\sqrt{\mu_{2p}}}{m+2}y^{\frac{m+2}{2}}\right)e^{\left[\mu_{kp} - \ln\sqrt{\alpha_1^2+\alpha_2^2} - i\left(\arctan\frac{\alpha_2}{\alpha_1}+s\pi\right)\right]t},$$

where $D_{kp} = A_k \cdot B_p \cdot C_{kp}$ are constants.

Remark 1. One can easily see that $\lambda_1 < 0$ in (17), which contradicts to condition $\mathrm{Re}\lambda = \lambda_1 \geq 0$ of the theorem 2.

Remark 2. The following problems can be studied by similar way. Instead of condition (6) we put conditions as follows:

Problem's name	P$_3$	P$_4$	P$_5$	P$_6$	P$_7$	P$_8$	P$_9$	P$_{10}$
S$_2$	u_x	u	u	u_x	u	u	u_x	u
S$_3$	u_y	u	u_y	u	u_y	u	u	u
S$_4$	u	u_x	u	u_x	u	u_x	u	u
S$_5$	u	u_y	u_y	u	u	u	u	u_y

3 Non-local problem for "forward-backward" parabolic equation with parameter.

In the domain $D = D_1 \cup D_2 \cup I_0,\ \ D_1 = \left\{(x,y) : -1 \leq x \leq 0,\ 0 \leq y \leq 1\right\},$

$I_0 = \left\{(x,y) :\ x = 0,\ 0 \leq y \leq 1\right\},\ \ D_2 = \left\{(x,y) : 0 \leq x \leq 1,\ 0 \leq y \leq 1\right\}$ let us consider equation

$$Lu = \lambda u, \tag{19}$$

where $\lambda \in R$, $Lu = u_{xx} - sign\left(x\right)u_y$.

The problem 3. To find a regular solution of the equation (19) from the class of functions $u\left(x,y\right) \in C\left(\overline{D}\right) \cap C^1\left(D \cup I_1 \cup I_2\right)$, satisfying non-local conditions

$$k_1u_x(-1,y) + k_2u(-1,y) = k_3u_x(1,y),\ \ 0 \leq y \leq 1, \tag{20}$$

$$k_4u_x(1,y) + k_5u(1,y) = k_6u_x(-1,y),\ \ 0 \leq y \leq 1; \tag{21}$$

$$u(x,0)=\alpha u(x,1), \quad -1\le x\le 1. \tag{22}$$

Here $k_i\ \left(i=\overline{1,6}\right)$, α is given non-zero constant, $I_1=\{(x,y):\ x=-1,\ 0\le y\le 1\}$, $I_2=\{(x,y):\ x=1,\ 0\le y\le 1\}$.

Note, non-local conditions (20), (21) were used for the first time by N.I.Ionkin and E.I.Moiseev [26, 27].

Theorem 3. If

$$|\alpha|=1,\ \lambda>0,\ k_3k_5=k_2k_6,\ k_1k_2<0,\ k_4k_5>0 \tag{23}$$

and exists a solution of the problem 3, then it is unique.

Proof:

We multiply equation (19) to the function $u(x,y)$ and integrate along the domains D_1 and D_2. Using Green's formula and condition (22), we get

$$\int_0^1 u(-0,y)\,u_x(-0,y)\,dy=\int_{-1}^0\frac{\alpha^2-1}{2}u^2(x,1)\,dx+\int_0^1 u(-1,y)\,u_x(-1,y)\,dy+\iint_{D_1}\left(u_x^2+\lambda u^2\right)dxdy,$$

$$\int_0^1 u(+0,y)\,u_x(+0,y)\,dy=\int_0^1\frac{\alpha^2-1}{2}u^2(x,1)\,dx+\int_0^1 u(1,y)\,u_x(1,y)\,dy-\iint_{D_2}\left(u_x^2+\lambda u^2\right)dxdy.$$

From conditions (20), (21), we find

$$u(1,y)\,u_x(1,y)=\frac{k_6}{k_5}u_x(-1,y)\,u_x(1,y)-\frac{k_4}{k_5}u_x^2(1,y),$$

$$u(-1,y)\,u_x(-1,y)=\frac{k_3}{k_2}u_x(-1,y)\,u_x(1,y)-\frac{k_1}{k_2}u_x^2(-1,y).$$

Taking ainto account above identities we establish

$$\begin{gathered}\int_{-1}^0\tfrac{\alpha^2-1}{2}u^2(x,1)\,dx+\int_0^1\tfrac{1-\alpha^2}{2}u^2(x,1)\,dx+\int_0^1\left[\tfrac{k_4}{k_5}u_x^2(1,y)-\tfrac{k_1}{k_2}u_x^2(-1,y)\right]dy+\\+\int_0^1\left[\tfrac{k_3}{k_2}-\tfrac{k_6}{k_5}\right]u(-1,y)\,u_x(1,y)\,dy+\iint_{D_1}\left(u_x^2+\lambda u^2\right)dxdy+\iint_{D_2}\left(u_x^2+\lambda u^2\right)dxdy.\end{gathered}$$

Considering condition (23), we get $u(x,y)\equiv 0$ in D an the proof of theorem is complete.

Remark 3. By similar method one can prove the uniqueness of solution of boundary-value problem with non-local initial condition for equation

$$0=\begin{cases} y^m u_{xx}+(-x)^n u_y-\lambda(-x)^n y^m u=0, & x<0\\ y^m u_{xx}-x^n u_y-\lambda x^n y^m u=0, & x>0.\end{cases}$$

Open question. A question is still open, on the unique solvability of boundary value problems for the following equation:

$$0=\begin{cases} y^{m_1}(-x)^{n_2}u_{xx}+(-x)^{n_1}y^{m_2}u_y-\lambda_1 u=0, & x<0\\ y^{m_1}x^{n_2}u_{xx}-x^{n_1}y^{m_2}u_y-\lambda_2 u=0, & x>0,\end{cases}$$

where λ_1, λ_2 are given complex numbers and $m_i,\ n_i=const>0\ \ (i=1,2)$.

References

1. Frankl F.I. On the problems of Chaplygin for mixed subsonic and supersonic flows. Izv.Akad. Nauk SSSR Ser. Mat., 1945, No 9, pp. 121-143.
2. Smirnov M.M. Degenerate elliptic and hyperbolic equations. Moscow: Nauka, 1966, 292 p.
3. J.M.Rassias, Uniqueness of Quasi-Regular Solutions for a Bi-Parabolic Elliptic Bi-Hyperbolic Tricomi Problem. Complex Variables, 2002, Vol.47, No 8, pp.707-718.
4. J.M.Rassias, Mixed Type Partial Differential Equations in R^n, Ph.D. Dissertation, university of California, Berkeley, USA, 1977, 1-135.
5. J.M.Rassias, Lecture Notes on Mixed Type Partial Differential Equations, World Scientific, 1990, pp.1-144.
6. J.M.Rassias, Mixed type partial differential equations with initial and boundary values in fluid mechanics. Int.J.Appl.Math.Stat. 13(2008), No J08, 77-107.
7. J.M.Rassias and A.Hasanov, Fundamental solutions of two degenerated elliptic equations and solutions of boundary value problems in infinite area. Int.J.Appl.Math.Stat. 8(2007), No M07. 87-95.
8. J.M.Rassias, Tricomi-Protter problem of nD mixed type equations. Int.J.Appl.Math.Stat. 8(2007), No M07. 76-86.
9. J.M.Rassias, Existence of weak solutions for a parabolic elliptic-hyperbolic Tricomi problem. Tsukuba J.Math., 23(1999), no.1. 37-54.
10. J.M.Rassias, Uniqueness of quasi-regular solutions for a parabolic elliptic-hyperbolic Tricomi problem. Bull.Inst.Math.Acad.Sinica, 25(1997), no. 4. 277-287.
11. G.C.Wen, The Exterior Tricomi Problem for Generalized Mixed Equations with Parabolic Degeneracy. Acta Mathematics Sinica, English Series, Vol.22, No 5, pp. 1385-1398.
12. G.C.Wen and D.Chen, Discontinuous Riemann-Hilbert problems for quasilinear degenerate elliptic complex equations of first order. Complex Var. Theory Appl. 50(2005), no. 7-11. 707-718.
13. G.C.Wen, The mixed boundary-value problem for second order elliptic equations with degenerate curve on the sides of an angle. Math. Nachr. 279(2006), no.13-14. 1602-1613.
14. S.Huang, Y.Y.Qiao and G.C.Wen, Real and complex Clifford analysis. Advances in Complex Analysis and its Applications, 5. Springer, New York, 2006. x+251.
15. G.C.Wen and H.G.W.Begehr, Boundary value problems for elliptic equations and systems. Pitman Monographs and Surveys in Pure and Applied Mathematics, 46. Longman Scientific and Tech., Harlow; John Wiley and Sons, Inc.,N.Y., 1990. xii+411.
16. A.Hasanov, Fundamental solutions of generalized bi-axially symmetric Helmholtz equation. Complex Variables, 2007, Vol. 52, No. 8, pp. 673-683.
17. M.Gevrey, Sur les equations aux derivees partielles du type parabolique. J.Math.Appl., 1914, Ch.4, pp.105-137.
18. A.Friedman, Fundamental solutions for degenerate parabolic equations. Acta Mathematica 133. Imprime le 18 Fevrier 1975, pp. 171-217.

19. Yu.P.Gorkov, Construction of a fundamental solution of parabolic equation with degeneration. Calcul. methods and programming, 2005, V.6, pp.66-70.

20. N.N.Shopolov, Mixed problem with non-local initial condition for a heat conduction equation. Reports of Bulgarian Academy of Sciences, 1981, V.34, No 7, pp.935-936.

21. A.A.Kerefov, On a spectra of the Gevrey problem for loaded mixed parabolic equation. Proceeding of the conference:Classical and non-classical boundary-value problems for partial differential equations, special functions, integral equations and their applications, 25-29 April, Russia, pp.76-77.

22. K.B.Sabitov, To the theory of mixed parabolic-hyperbolic type equations with spectral parameter. Differencial'nye Uravnenija, 1989, Vol.25, No 1, pp. 117-126.

23. A.S.Berdyshev and E.T.Karimov, Some non-local problems for the parabolic-hyperbolic type equation with non-characteristic line of changing type. CEJM, 2006, Vol. 4, No 2, pp. 183-193.

24. E.T.Karimov, About the Tricomi problem for the mixed parabolic-hyperbolic type equation with complex spectral parameter. Complex Variables, 2005, Vol.56, No 6, pp.433-440.

25. E.T.Karimov, Some non-local problems for the parabolic-hyperbolic type equation with complex spectral parameter. Mathematische Nachrichten, Vol.281, Iss. 7, 2008, pp.959-970.

26. N.I.Ionkin, The stability of a problem in the theory of heat condition with non-classical boundary conditions. (Russian). Differencial'nye Uravnenija 15 (1979), No 7, 1279-1283, 1343.

27. N.I.Ionkin and E.I.Moiseev, A problem for a heat equation with two-point boundary conditions. (Russian). Differencial'nye Uravnenija 15 (1979), No 7, 1284-1295, 1343.

Name: JOHN MICHAEL RASSIAS
Affiliation: Professor
Institute: National and Capodistrian University of Athens
Postal address: 4, Agamemnonos Str., Aghia Paraskevi, Athens 15342, Greece

Name: KARIMOV ERKINJON TULKINOVICH
Affiliation: Junior Scientific Researcher
Institute: Institute of Mathematics and information technologies, Uzbek Academy of Sciences
Postal address: 29, Durmon yuli str., Tashkent-100125, Uzbekistan.